

2022 Fact Sheet

on Women and Men in the Philippines



Health and Nutrition				
Indicator	Women	Men	Ref. period	Source
Proportion of Underweight Children 0-5 Years Old (0-60months) a/	19.3	18.8	2018-2019	ENNS, FNRI
Proportion of Underweight Children 5.08-10 Years Old (61-120 months) a/	22.9	28.0		
Proportion of Stunted Children 0-5 Years Old (0-60months) b/	28.7	30.3		
Proportion of Stunted Children 5.08-10 Years Old (61-120 months) b/	23.1	26.5		
Proportion of Overweight for Height Children 0-5 Years Old (0-60months) c/	3.1	3.9		
Proportion of Overweight/Obese Children 5.08-10 Years Old d/	9.4	11.5		
Proportion of Obese Among Adults 20 Years and Above e/	11.1	6.1		
Proportion of Wasting Children less than 0 - 5 Years Old (0 - 60 months) c/	5.3	6.4		
Proportion of Wasting Children less than 5.8 - 10 Years Old (61 - 120 months) c/	7.1	8.5		_
Percentage of Children 12-23 Months Old Who Received All Basic Vaccinations at Anytime Before ne Survey (%) f/	70.8	69.2	2017	NDHS, PSA
Neonatal Mortality Rate (per 1,000 live births) for the 5-year period preceding the survey	14.0	13.0		
Postneonatal Mortality Rate (per 1,000 live births) for the 5-year period preceding the survey	14.0	10.0		
Sourceonatal Mortality Rate (per 1,000 live birtils) for the 3-year period preceding the survey	70	8.0		
a fact Mantality Data (and 1000 live birthe) factors for an and a more directly a survey	7.0			
nfant Mortality Rate (per 1,000 live births) for the 5-year period preceding the survey Child Mortality Rate (per 1,000 live births) for the 5-year period preceding the survey	6.0	21.0 7.0		
Jnder-Five Mortality Rate (per 1,000 live births) for the 5-year period preceding the survey	26.0	28.0		
	28.0	28.0		
Percentage of Currently Married Women Aged 15-49 Years Using Contraception (%)	54.3	-		
Percentage of Women Aged 15 - 49 Years With Unmet Need for Family Planning				
Currently Married Women	17.0	-		
Sexually Active Unmarried Women	49.0	-		
Proportion of Births in the 5 Years Preceeding the Survey Attended by Skilled Health Professional	84.4	_		
Top Leading Causes of Mortality Rate (per 100,000 persons)				
Diseases of Circulatory System	91,184	122,441	2019	DOH
Diseases of the Respiratory System	41,876	54,003		
Neoplasms	36,115	32,542		
Adolescent Birth Rates per 1,000 Women Aged 15-19 Years	47.0	-	2017	NDHS, PSA
Percentage of Women Who Are Pregnant and Who Have Begun Child Bearing at Age 15-19 Years	9.0	-		
Number of Newly-Diagnosed HIV Cases, by Sex and Age (as of Dec. 2021)	7.0	-	2021	HARP, DOH
Monthly (Dec 2021)	41	952	2021	
Annual (Jan to Dec 2021)	548	11,793		
Cumulative (Jan 1984 - Dec 2021)	5,508	88.818		
	5,508	88,818		

Notes:

* - indicator(s) is/are part of the updated Philippine Core GAD Indicators. More information on the Updated Philippine Core GAD is accessible through this link https://psa.gov.ph/psa-board-4

-- not applicable or data not reported

a/ Weight-for-age classification based on WHO-Child Growth Standards

b/ Height-for-age classification based on WHO-Child Growth Standards

c/ Weight-for-height classification based on WHO-Child Growth Standards

d/ Overweight and obese based on Body Mass Index - for - age classification from WHO Grownth References 2007

e/ Covers adults aged 20 years old and over and based on World Health Organization - Body Mass Index

f/ It is based on the vaccination card or the mother's report. All vaccinations include Bacillus, Calmette-Guerin (BCG), measles, three doses each of the diphtheria, pertussis, tetanus (DPT),

polio and Hepatitis- B vaccine (either Hepa-B0, B1 and B2 or Hepa-B1, B2 and B3); excludes Haemophilus influenzae type B (HiB) vaccine.

Data Source:

ENNS - Expanded National Nutrition Survey NDHS - National Demographic and Health Survey HARP - HIV/AIDS & Art Registry of the Philippines

Data Agencies:

DOH - Department of Health

FNRI - Food and Nutrition Research Institute PSA - Philippine Statistics Authority



Alternation and a second